

TRIAL SUMMARY

Crop Type: Corn Silage

Year: 2021

Location: Darcy Frankl, Raymore, SK

CANTERRA SEEDS Contact: Bret Gaetz

Planting Date: May 28, 2021

Harvest Date: Sept 30, 2021

Trial Type: 1/1000 Scale

Row Width: 30 inches

PRIDE Seed Contact: Sara Meidlinger



COMPANY	VARIETY	CHU	RM	Moisture %	DM %	Protein %	ADF %	NDF %	STARCH %	TDN %	NE/g	NE/l	TONS/AC ACTUAL	TONS/AC AT 65%	DRY Yield RANK	MILK LB/AC	MILK RANK	BEEF LB/AC	BEEF RANK
Pioneer	P7005AM	2000	70	44.5	55.5	6.9	24	43.2	38.3	63	0.77	1.5	9.8	15.53	1	15,764	1	1142	1
Pioneer	P7211AM	2050	72	56.7	43.3	6.3	26.8	48.1	32.8	60	0.67	1.38	9.7	12.01	7	10,872	9	840	8
Northstar	913S	2105	75	61.5	38.5	7	24.6	45.7	32.2	62	0.73	1.45	11.7	12.81	4	13,997	5	926	4
Northstar	932S	2400	89	76.9	23.1	9.8	29.9	54.8	15.6	59	0.61	1.25	13.6	8.95	11	8,786	11	616	11
Northstar	NS271	2050	71	68.9	31.1	10.3	25.5	47.6	24.4	60	0.68	1.38	10.4	9.19	10	9,946	10	643	10
Pickseed	PS2333RR	2275	77	73.7	26.3	9.1	22.6	45.9	27.9	62	0.74	1.44	17.9	13.44	2	14,670	2	972	2
Pickseed	PS2320RR	2200	76	68.8	31.2	9.7	24.7	44.3	26.2	62	0.76	1.46	14.2	12.68	5	14,152	4	917	5
PRIDE Seeds	A4646G2 RIB	2300	79	69.9	30.1	9	25.4	47.4	23.9	59	0.63	1.34	14.4	12.37	6	12,506	7	852	6
PRIDE Seeds	AS1017RR EDF	2200	74	71.5	28.5	9.3	27.9	48	23.5	60	0.67	1.36	14.6	11.86	8	12,260	8	830	9
PRIDE Seeds	AS1027RR EDF	2425	80	69.7	30.3	8.4	23.3	43	29.9	61	0.72	1.45	15.2	13.18	3	14,242	3	938	3
PRIDE Seeds	A4414RR	2150	74	69.6	30.4	9.3	24.2	47.5	25	61	0.69	1.39	13.7	11.85	9	12,541	6	843	7

Nutrient	Target Value	Definitions	Reasoning
Dry Matter (DM)	30-40%	The percentage of forage that is not water	Excessive moisture content can cause spoilage and decrease silage quality. Too dry is usually associated with reduced digestibility and energy content.
Crude Protein (CP)	7-9%	Total amount of nitrogen (N) in a forage.	High protein is desirable. Low protein may be caused by under fertilization, nitrogen competition, or improper harvesting and/or storage.
Acid Detergent Fiber (ADF)	20-33%	Percent of highly indigestible material in a forage. Comprised of cellulose, lignin, cutin, silica, pectin, and unavailable protein.	High ADF content is an issue for the same reasons as high NDF content. ADF is negatively correlated to digestibility and energy
Neutral Detergent Fiber (NDF)	35-55%	Partially available to animals. Percent of cell wall material in a forage; cellulose, hemicelluloses, Lignin, cutin, and unavailable protein.	NDF values will generally increase with low grain silage, stress, or immaturity. NDF is an inverse predictor of intake. (higher NDF equals lower intake and visa versa)
Starch	>28%	Form of carbohydrates stored in plants. It is the specific polysaccharide of many glucose subunits.	Usually higher content is better. If starch levels are <28% this usually indicates the silage was cut early or the crop was stressed.
Total Digestible Nutrients (TDN)	67-74%	Sum of all digestible organic nutrients that are available to the animal, as a % or DM.	Could be used to express the energy value of the corn silage.
Net Energy for Lactation (NEl)	>0.64% Mcal/lb	An estimate of the energy value of a feed used for milk production	Mega calories of energy for lactation. Higher values usually indicate a better-quality corn silage.
Net Energy for Gain (NEg)	0.4-0.5 Mcal/lb	An estimate of the energy for weight gain. Energy above maintenance.	Mega calories of energy for gain.