

# TRIAL SUMMARY

**Crop Type:** Corn Silage  
**Year:** 2021  
**Location:** Trehane Holsteins, Niverville, MB  
**Retail:** GJ Chemical – Arnaud, MB  
**CANTERRA SEEDS Contact:** Jackie Dudgeon

**Planting Date:** April 30, 2021  
**Harvest Date:** Sept 17, 2021  
**Trial Type:** Field Scale  
**Row Width:** 20 inches  
**PRIDE Seed Contact:** Sara Meidlinger



COMPANY	VARIETY	CHU	RM	Moisture %	DM %	Protein %	ADF %	NDF %	STARCH %	TDN %	NE/g	NE/l	TONS/AC ACTUAL	TONS/AC AT 65%	DRY Yield RANK	MILK LB/AC	MILK RANK	BEEF LB/AC	BEEF RANK
Dekalb	DKC 35-37 RIB	2575	85	69.4	30.6	7.7	22.2	41.8	32.8	64	0.83	1.54	15.9	13.94	6	15,582	7	1041	6
Dekalb	DKC 36-86 RIB	2625	86	68.4	31.6	8.4	21.9	38.9	35	64	0.82	1.56	16.7	15.12	5	16,480	5	1129	5
Dekalb	DKC 33-37 RIB	2500	83	67.9	32.1	8.6	19.6	36.1	36.7	65	0.86	1.60	19.9	18.29	3	20,765	3	1387	3
<b>PRIDE</b>	<b>AS1047RR EDF</b>	<b>2450</b>	<b>81</b>	<b>68.7</b>	<b>31.3</b>	<b>8.6</b>	<b>23.9</b>	<b>42</b>	<b>30</b>	<b>65</b>	<b>0.84</b>	<b>1.54</b>	<b>23.2</b>	<b>20.79</b>	<b>1</b>	<b>23,080</b>	<b>1</b>	<b>1576</b>	<b>1</b>
Thunder	TH6185 VT2P	2550	85	67.1	32.9	8.4	18.3	33.9	38.6	66	0.92	1.66	14.1	13.25	7	15,640	6	1020	7
Thunder	TH6183 VT2P	2450	83	67.4	32.6	8	21.2	38.7	30.7	62	0.77	1.51	14.1	13.13	8	14,371	8	950	8
Pioneer	P8820AM	2600	88	71.3	28.7	8.9	20	36.8	34.3	64	0.85	1.59	23.2	19.02	2	21,989	2	1420	2
Maizex	Venza R	2600	88	73.9	26.1	8.6	22.7	43	29.5	62	0.75	1.47	23.0	17.15	4	19,303	4	1240	4

Nutrient	Target Value	Definitions	Reasoning
Dry Matter (DM)	30-40%	The percentage of forage that is not water	Excessive moisture content can cause spoilage and decrease silage quality. Too dry is usually associated with reduced digestibility and energy content.
Crude Protein (CP)	7-9%	Total amount of nitrogen (N) in a forage.	High protein is desirable. Low protein may be caused by under fertilization, nitrogen competition, or improper harvesting and/or storage.
Acid Detergent Fiber (ADF)	20-33%	Percent of highly indigestible material in a forage. Comprised of cellulose, lignin, cutin, silica, pectin, and unavailable protein.	High ADF content is an issue for the same reasons as high NDF content. ADF is negatively correlated to digestibility and energy
Neutral Detergent Fiber (NDF)	35-55%	Partially available to animals. Percent of cell wall material in a forage; cellulose, hemicelluloses, Lignin, cutin, and unavailable protein.	NDF values will generally increase with low grain silage, stress, or immaturity. NDF is an inverse predictor of intake. (higher NDF equals lower intake and visa versa)
Starch	>28%	Form of carbohydrates stored in plants. It is the specific polysaccharide of many glucose subunits.	Usually higher content is better. If starch levels are <28% this usually indicates the silage was cut early or the crop was stressed.
Total Digestible Nutrients (TDN)	67-74%	Sum of all digestible organic nutrients that are available to the animal, as a % or DM.	Could be used to express the energy value of the corn silage.
Net Energy for Lactation (NEl)	>0.64% Mcal/lb	An estimate of the energy value of a feed used for milk production	Mega calories of energy for lactation. Higher values usually indicate a better-quality corn silage.
Net Energy for Gain (NEg)	0.4-0.5 Mcal/lb	An estimate of the energy for weight gain. Energy above maintenance.	Mega calories of energy for gain.