

**Varietal Characteristics:**

- **Growing Zone:** Suited to all pea-growing zones.
- **Yield:** Excellent
- **Maturity:** Earlier than CDC Amarillo
- **Vine length:** Longer than Agassiz
- **Lodging:** Better than Agassiz
- **Seed Size:** Intermediate
- **Seed Coat Breakage:** Better than CDC Amarillo
- **Disease Resistance:** R to powdery mildew, MS to Fusarium Wilt, MS to Mycosphaerella Blight

**Technical Information:**

**2018 Seed Manitoba**

Variety	Yield % CDC Meadow	Site Years Tested	Relative Maturity	Seed Size	Lodging	Powdery Mildew	Seedcoat Breakage (%)	Relative Vine Length
<b>AAC Carver</b>	<b>110</b>	<b>10</b>	<b>M</b>	<b>240</b>	<b>G</b>	<b>VG</b>	<b>G</b>	<b>L</b>
AC™ Agassiz	101	53	M	230	G	VG	G	M
CDC Amarillo	103	16	M	230	VG	VG	F	M
CDC Meadow	100	66	E	220	G	VG	G	M

**2018 SaskSeed Guide**

Variety	Yield - % CDC Amarillo			Yrs Tested	Rel Mat	Lodging (1-9)	Seed Coat Dimpling	Seed Weight (g/1000)	Protein vs CDC Amarillo
	1,2 & South 3	North 3 & 4	Irrigation						
<b>AAC Carver</b>	<b>103</b>	<b>101</b>	--	<b>4</b>	<b>E</b>	<b>4.0</b>	<b>F</b>	<b>240</b>	<b>-1.6</b>
AC™ Agassiz	98	93	100	9	M	4.5	F	230	0.3
CDC Amarillo	100	100	100	9	M	3.5	F	230	23.0
CDC Meadow	91	89	90	9	E	4.0	G	220	-0.6

**2018 Alberta Seed Guide**

Variety	Yield % CDC Amarillo	Overall Station yrs	Maturity Rating	Vine Length (cm)	TSW (g)	Lodging (1-9)	Seed Coat Breakage	Seed Coat Dimpling
<b>AAC Carver</b>	<b>106+</b>	<b>28</b>	<b>E</b>	<b>86</b>	<b>240</b>	<b>3.4</b>	<b>G</b>	<b>G</b>
CDC Amarillo	100	91	M	80	226	2.6	F	F
CDC Meadow	96-	77	M	81	203	3.9	G	G
AAC Lacombe	104+	61	M	75	256	2.4	G	F

**Origin:** AAFC, Lacombe Research Station (Test Code: MP 1920)

**Pedigree:** CDC715S-4/P98089063